


AD-A151 959

AFOSR-TR

WJSA R85T-03

②



SHORT WAVELENGTH CHEMICAL LASER (SWCL) WORKSHOP

Interim Report under contract F49620-84-C-0104

5 December 1984

Submitted to:
Air Force Office of Scientific Research
Bolling AFB
Washington, DC 20332

and

Strategic Defense Initiative Organization
Office of the Secretary of Defense
Washington, DC 20301

DTIC FILE COPY

Prepared by:
W. J. Schafer Associates, Inc.
1901 N. Fort Myer Drive, Suite 800
Arlington, Va 22209

85 03 13 017

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution Unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) WJSA R85T-03			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 85-0277		
5a. NAME OF PERFORMING ORGANIZATION W. J. Schafer, Assocs Inc		5b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NP		
6a. ADDRESS (City, State and ZIP Code) 1901 N. Fort Myer Drive, Suite 800 Arlington, VA 22209			7b. ADDRESS (City, State and ZIP Code) Building 410 Bolling AFB DC 20332-6448		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable) NP	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620-84-C-0104		
8c. ADDRESS (City, State and ZIP Code) Building 410 Bolling AFB DC 20332-6448			10. SOURCE OF FUNDING NOS		
			PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2301	TASK NO. A1
11. TITLE (Include Security Classification) SHORT WAVELENGTH CHEMICAL LASER (SWCL) WORKSHOP					
12. PERSONAL AUTHOR(S) Dr William Watt					
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 1 Sep 84 TO 31 Aug 85		14. DATE OF REPORT (Yr., Mo., Day) 5 Dec 84	
15. PAGE COUNT 40					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The workshop was held for the purpose of identifying the government's interest in SWCL technology, reviewing past and present efforts in this area and presenting the government's plans for a new thrust in SWCL source development. In addition, the workshop was to provide a forum for interaction between members of the Strategic Defense Initiation Organization (SDIO) and the 6.1 agencies with the technical community in order to create an enthusiastic response to the SWCL thrust and to generate new concepts as well as to involve new participants in this technically challenging area.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL DR HOWARD R. SCHLOSSBERG		22b. TELEPHONE NUMBER (Include Area Code) 202/767-4906		22c. OFFICE SYMBOL NP	

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

SHORT WAVELENGTH CHEMICAL LASER (SWCL) WORKSHOP

Interim Report under contract F49620-84-C-0104

5 December 1984

Submitted to:
Air Force Office of Scientific Research
Bolling AFB
Washington, DC 20332

and

Strategic Defense Initiative Organization
Office of the Secretary of Defense
Washington, DC 20301

A-1

Prepared by:
W. J. Schafer Associates, Inc.
1901 N. Fort Myer Drive, Suite 800
Arlington, Va 22209

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)

W. J. Schafer Associates, Inc. (WJSA) is pleased to submit this summary report on the Short Wavelength Chemical Laser (SWCL) workshop held 14, 15 November 1984 in Charleston, S.C. Attached to this report are the workshop program, the attendance list and abstracts of the technical papers presented.

The workshop was held for the purpose of identifying the government's interest in SWCL technology, reviewing past and present efforts in this area and presenting the government's plans for a new thrust in SWCL source development. In addition, the workshop was to provide a forum for interaction between members of the Strategic Defense Initiative Organization (SDIO) and the 6.1 agencies with the technical community in order to create an enthusiastic response to the SWCL thrust and to generate new concepts as well as to involve new participants in this technically challenging area.

By any of these standards the SWCL workshop was a success. Furthermore, WJSA believes, as a result of extremely favorable comments and letters from a large number of attendees, that the workshop was exceptionally well organized and thereby provided the appropriate environment for achieving the desired objectives.

The interest of the technical community in the SWCL thrust was demonstrated by the 142 attendees from academic, government laboratory, industry and national laboratory research groups. The attendance, which was larger than anticipated, included experts in all the scientific disciplines required for the development of high-power laser sources. Further, as a result of the enthusiastic response of the technical community, the workshop sessions were extended to include papers which were submitted voluntarily. While this restricted somewhat the opportunities for open session discussions it generated a significant level of interaction behind the scenes as well as an exceptional attendance level at all of the technical sessions.

Dr. L. Marquet of SDIO opened the workshop by reviewing the philosophy behind the strategic defense initiative research effort and outlining the goals for the SWCL thrust. He introduced the key SDIO personnel and the 6.1 Agencies through which the research efforts would be contracted and identified the format and schedule for preparing responses. These remarks set the stage well for the review talks which followed immediately and for the more detailed technical sessions.

Since one purpose of the workshop was to inject new blood and concepts into the area, the technical program was designed to provide both a comprehensive review as well as information to guide future efforts. Both objectives were more than satisfied. All current efforts were described from which a consensus was reached, that energy transfer processes are to be preferred over direct excitation for the development of chemically pumped short wavelength lasers. In this regard the organization of the workshop to include sessions on the generation of metastable excited species and on the energy transfer of these and other species was well conceived and extremely valuable.

The overview session was highlighted by Dr. Huestis' talk which laid out the electronic excitation pathways that are required for SWCL development. Other talks described what has been learned from the HF and COIL laser efforts.

In addition to the Energy Transfer session which was mentioned above, the technical presentations were grouped as follows: Metal/Oxidizer Systems; Pyrotechnic Systems; Metastable Species Production and Metastable Transfer Systems. The program which gives the titles and authors of each paper and the names of the session chairs is attached.

Each session opened with a review paper in which the general status of the technology was described by a well known authority in that field. These were followed by more detailed discussions of particular chemi-excitation mechanisms and approaches to short wavelength chemical lasers.

In the metal/oxydizer session interesting papers were given by Professors Djeu and Parsons. Djeu described how the BaO^*/Ba transfer had led to a no-loss/no-gain situation on the 1.3 micron transition in Ba while Parsons showed that oxidation reactions of metastable atoms provided higher excited state species densities than could be achieved with ground state atoms.

The use of chemi-excitation light sources for pumping Nd:YAG and other solid-state lasers was described by three authors. Although the results were as yet inconclusive for tactical military applications this approach was found to be more costly than flashlamp excitation for laser fusion.

The session on metastable state production was highlighted by an excellent introduction by Dr. Heidner of Aerospace. He proposed $NF(b)$ transfer to IF and $NF(a)$ transfer to BiF as potential SWCL candidates. Professor Coombe's examination of the potential of azide decomposition for the production of $N_2(A)$ species and Dr. Rossi's discussion of dioxetane decomposition to produce long lived excited ketones were two examples of novel excitation routes to SWCLs.

The IF system and other interhalogens were introduced by Dr. Davis of AFWL. IF appears to be a promising SWCL candidate and has been lased under both optical and electrical excitation of mixtures of iodine with fluorinated hydrocarbons. Davis projected that 10^{-3} - 10^2 torr of IF would almost certainly lase since vibrational relaxation would concentrate the upper level species in the lowest vibrational state and deactivate the lower level ($V'' = 4-7$) which exhibited a significant Frank-Condon shift. Subsequent speakers discussed the rates of energy transfer into IF(B) from $N_2(A)$, and those for intra- and intermolecular energy transfer processes.

The final session was concerned with the details of energy transfer processes and provided an interpretation of the atomic and molecular physics processes involved. The paper by Prof. George addressed atomic excitation in collisions between gas-phase species and surfaces which was very different than any other reactions that had been described.

In summary the workshop was well organized, well attended and provided a program which has provoked an enthusiastic response from the technical community. WJSA intends to keep current in ongoing and new activities in this area and suggest that the government plan a further workshop to review the status of technology in May 1986. WJSA believes that it has well demonstrated its capabilities to effectively plan and organize such a meeting and would be pleased to be selected again to perform these activities.

SHORT WAVELENGTH CHEMICAL LASER WORKSHOP

**MILLS HOUSE HOTEL
Charleston, S.C.**

13-15 November 1984

Final Program

November 14 - Hibernian Hall

- 8:15 - 8:45 Late Registration
- 8:45 - 8:55 Welcome & Administrative Remarks
Dr. W. Watt, WJSA
- 8:55 - 12:00 Plenary Session - Chair: Dr. L. Marquet, SDIO
- 8:55 - 9:10 Plan for SWCL Thrust
Dr. L. Marquet, SDIO
- 9:10 - 9:50 HF - What Have We Learned?
Dr. W. Warren, PAR
- 9:50 - 10:30 Chemical Oxygen - Iodine Laser Review
Dr. P. Avizonis, AFWL
- 10:30 - 10:45 COFFEE BREAK
- 10:45 - 11:25 Why So Few Chemical Lasers?
Dr. D. Huestis, SRI
- 11:25 - 12:00 Approach to Efficient Short-Wavelength
Chemical Lasers
Dr. P. Avizonis, AFWL
- 12:00 - 1:15 LUNCH
- 1:15 - 3:15 Metal/Oxidizer Systems
Chair: Dr. C. R. Jones, LANL
- 1:15 - 1:55 Potential of Metal/Oxidizer Systems
Prof. J. Gole, Georgia Tech
- 1:55 - 2:15 Review of BaO/Ba Transfer Laser Studies
Prof. N. Djou, Univ. of South Florida
- 2:15 - 2:35 SnO and SnF Studies
Dr. J. Raymonda, Bell Aerospace
- 2:35 - 2:55 Reactions of Transition and Main Group III
and IV Metals with Halogens
Prof. J. Parsons, Ohio State Univ.
- 2:55 - 3:15 Mg/N₂O/CO Flame Studies of Mg(3p) Production
Dr. R. Meinzer, UTRC
- 3:15 - 3:30 AFTERNOON BREAK

- 3:30 - 5:10 Pyrotechnic Systems - Chair: Dr. B. Feldman, NRL
- 3:30 - 4:10 Recent Developments in Chemiluminescence
Prof. A. Fontijn, RPI
- 4:10 - 4:30 Combustion - Pumped Solid State Lasers
Mr. E. Chicklis, Sanders
- 4:30 - 4:50 Chemical Flashlamps Employing F₂ Oxydizers
Dr. J. Waymouth, GTE
- 4:50 - 5:10 Luminous Metal Dust Oxide Reactions as
Potential Laser Pump Sources
Dr. M. Duignan, NRL

November 15 - Hibernian Hall

8:30 - 10:20 Metastable State Production
Chair: Dr. H. Powell, LLNL

- 8:30 - 9:00 Production and Uses of NF and N₂ Metastables
Dr. R. Hiedner, Aerospace
- 9:00 - 9:20 Excited-State Chemistry of Molecular Azides
Prof. R. Coombe, Univ. of Denver
- 9:20 - 9:40 Possible High Density Visible Chemical
Laser Based on Dioxetanes
Dr. M. Rossi, SRI
- 9:40 - 10:00 Chemiluminescence and Oxidation of Phosphorus
Prof. D. Stedman, Univ. of Denver
- 10:00 - 10:20 Excited Products from Extremely Exothermic
Reactions
Prof. C. Wittig, USC

10:20 - 10:30 COFFEE BREAK

10:30 - 12:30 Metastable Transfer Systems
Chair: Dr. S. Davis, AFWL

- 10:30 - 11:00 Overview of Interhalogen Lasers
Dr. S. Davis, AFWL
- 11:00 - 11:30 Electronic State Energy Transfer and Laser
Possibilities
Prof. D. Setser, Kansas State Univ.

- 11:30 - 11:50 $N_2(A) \rightarrow IF$ Transfer
Dr. L. Piper, PSI
- 11:50 - 12:10 $NF(b) \rightarrow IF$ Transfer
Dr. D. Benard, Rockwell Science Center
- 12:10 - 12:30 The NO ($A \rightarrow X$) Laser
Dr. M. Burrows, Los Alamos
- 12:30 - 1:45 LUNCH
- 1:45 - 3:40 Energy Exchange Mechanisms
Chair: Dr. M. Lavan, BMDATC
- 1:45 - 2:20 Vibration to Electronic Energy Transfer in
V-V Pumped Molecules
Dr. J. W. Rich, CALSPAN
- 2:20 - 2:40 Electronic Energy Transfer in Metal Atoms
Dr. S. Leone, JILA
- 2:40 - 3:00 Energy Transfer Processes Involving Atoms
and Molecules
Prof. J. Wiesenfeld, Cornell
- 3:00 - 3:20 Production of Excited Atomic States by
Ion-Surface Charge Exchange
Prof. T. George, Univ. of Rochester
- 3:20 - 3:40 Reactive and Inelastic Collisions of
Electronically Excited Species
Prof. M. Alexander, Univ. of Maryland
- 3:40 - 3:45 Review of Procedure and Schedule for Submitting
Proposals
- 3:45 ADJOURN

ABSTRACTS

Curt Wittig


Short Wavelength Chemical Laser Workshop

ABSTRACT

There are many chemical reactions involving atoms and small molecules, which are highly exoergic and do not involve particularly exotic substances. With some of these reactions, there is reason to believe that the products will be formed in excited states, and several such systems will be discussed in which the main products are N_2 and/or CO , the two strongest bonds formed in nature. We will use spin conservation and tight 4-center transition states in order to channel the enthalpy release into product internal states. Reactions are identified which are $\sim 1000 \text{ kJ mol}^{-1}$ exoergic and can produce metastable electronic states, and experiments will be described which measure nascent product distributions from these reactions.

CHEMILUMINESCENCE AND THE OXIDATION OF PHOSPHORUS. D.H. Stedman.
Chemistry Department, University of Denver, Denver, CO 80208

The oxidation of phosphorus and phosphine by various oxidants leads to a bright continuum emission extending from 600 to 800 nm. A quantum yield as high as 0.82 photons per PH_3 has been measured. These and other theoretical, mass spectroscopic and microwave studies combine to yield a suggested mechanism for the emission as



Evidence in favor of this mechanism and an effect of oxygen caused by the unique reaction $\text{PO} + \text{O}_2 \rightarrow \text{O} + \text{OPO}$ is presented, together with some spectroscopic speculation.


POSSIBLE HIGH DENSITY VISIBLE CHEMICAL LASER
BASED ON DIOXETANE

Michel J. Rossi
Chemical Physics Laboratory
337 International, Menlo Park, CA 94025

Thermal decomposition of substituted dioxetanes has been considered by many investigators as a possible approach to a visible chemical laser due to the high energy release and high yields of electronically excited products. However, the prospects for a gas-phase dioxetane chemical laser were judged slim because (1) the temperatures for rapid unimolecular decomposition could not be achieved fast enough and (2) the radiative properties of the excited state (triplet-excited carbonyl compounds) were not suitable for the upper laser level.

We report results of a modeling study where neat crystalline tetramethyldioxetane (TMDX) undergoes decomposition into acetone (A) triggered by irradiation in the near-UV. The salient features of our reaction mechanism are:

- The operation of a quantum chain reaction ($A = \text{acetone}$)



which heats up the sample with no loss in triplets. This reaction goes on until the temperature is high enough for the unimolecular decomposition to take over.

- Triplet-triplet energy pooling that results in production of singlet-acetone, which is proposed as the upper laser level:



We will present detailed results on the kinetic modeling in terms of initiation conditions, incubation times, and profiles of concentrations, temperatures, and pressures versus time. We will use these modeling studies to discuss the potential of this explosive reaction system as a chemical laser.

EXCITED STATE CHEMISTRY
OF MOLECULAR AZIDES

Robert D. Coombe

Department of Chemistry, University of Denver
Denver, Colorado 80208

Molecular azides have been shown to be uniquely capable of producing electronically excited species upon photodissociation or reaction with atoms or free radicals. For example, the halogen azides (XN_3 , $\text{X} = \text{halogen}$) are metastable with respect to dissociation to nitrogen halide diatomics and N_2 . In the photodissociation of these compounds, the energy available to the fragments is the sum of the photon energy and the energy borne by the azide, such that very highly excited states are accessible. In this regard, pulsed laser photodissociation of ClN_3 and BrN_3 has been shown to provide a unique source of $\text{N}_2(\text{A}^3\Sigma_u^+)$ metastables, which can undergo reaction or energy transfer processes with other species present. Reactions of molecular azides with atoms (e.g., halogens, nitrogen, and oxygen) have been shown to efficiently produce electronically excited nitrogen halides, N_2 and NO . The high specificity observed in these processes is largely a consequence of the chemistry of N_3 radicals, which act as an intermediate. The properties of these systems suggest their potential utility in a number of possible chemical laser systems.

CHEMICAL PRODUCTION OF METASTABLE NF AND N₂

R. F. Heidner III
Aerophysics Laboratory
The Aerospace Corporation

ABSTRACT

Our laboratory has endeavored to produce high densities of electronic metastables in order to exploit the energy storage potential of such systems for the production of chemically pumped electronic transition lasers. Early studies of Herbelin et al, firmly established the efficiency of the $H + NF_2$ reaction for producing $NF(a^1\Delta)$ and correctly predicted the reaction pathways leading to the production of $N(^2D)$ and N_2^* under H atom rich conditions.

Subsequently, studies were performed to scale the production of $NF(a)$ and $NF(b)$ and to utilize these species both for direct lasing experiments and for transfer laser concepts to suitable atomic and molecular receptors. The presentation will be both a review and a status report on current theory and experiments.

This work has been sponsored by the Air Force Weapons Laboratory under U.S. Air Force Space Division (AFSD) Contract FO4701-84-C-0085 and by the Aerospace Corporation (Aerospace Sponsored Research and Mission Oriented Investigation and Experimentation).

Luminous Metal Dust/Oxidation Reactions
as Potential Laser Pump Sources

Laser Physics Branch,
Naval Research Laboratory
Washington, DC 20375

Michael T. Duignan,* R.W. Waynant, and B.J. Feldman

Highly luminous metal oxidation reactions are a potentially efficient pump source for IR and near IR lasers. Metal powders or dusts offer unique advantages for repetitively pulsed or quasi-continuous pumping in that they are easily entrained in a gas and flowed.

We have conducted preliminary experiments with aluminum, magnesium, and zirconium powders in oxygen atmospheres. Optical emission as a function of time, time-integrated spectra, and dynamic pressure were monitored.

The potential for pumping solid state lasers with metal dust oxidation reactions will be discussed.

* Potomac Photonics, Inc., Alexandria, VA 22303

October 1984

RECENT DEVELOPMENTS IN CHEMILUMINESCENCE

Arthur Fontijn

Rensselaer Polytechnic Institute
Department of Chemical Engineering & Environmental Engineering
Troy, NY 12180-3590

In August a Symposium was held at the A.C.S. National Meeting on "Gas-Phase Chemiluminescence and Chemi-Ionization". Most of the invited presentations there plus a few additional papers, will be published in a book by the same title as the symposium (North-Holland, May 1985, A. Fontijn, Ed.). In this talk some of the presentations will be reviewed, with emphasis on points of interest to SWCL development. Subjects covered will include selective reaction by particular spin-orbit states, non-statistical population of excited product states, extraction of initial population information from chemiluminescent spectra, chemiluminescent chemi-ionization, chemiluminescence from ion-molecule reactions and the heterogeneous production of excited states. Earlier observations on the highly efficient (high quantum yield ϕ) $\text{Sn}/\text{N}_2\text{O}$ reaction will be reviewed. These observations demonstrate the importance of taking reaction temperature and activation energy into account when selecting reactions for laser pumping. A new approach to a high temperature SWCL will be suggested. It is important to realize that a low ϕ reaction can nonetheless have a large branching ratio R^* for excited state production, which (rather than ϕ) is the significant factor in selecting pumping reactions. To determine R^* , radiative lifetime and quenching rate coefficients have to be considered. Results from a novel method for obtaining this information for longer-lived electronically excited species will be shown.

Mg/N₂O/CO Flame Studies of Mg [³P] Production

R. A. Meinzer, H. H. Michels and R. Tripodi
United Technologies Research Center
East Hartford, CT 06108

Metastable magnesium atoms, [Mg³P], appear to represent an attractive energy storage candidate for powering a chemically pumped electronic transition energy transfer laser. They have a relatively long radiative lifetime, 4.5 ms; deactivation of Mg[³P] by various reagents is slow; spontaneous emission from the metastable electronic energy level to the ground state is in the blue region, ³P-¹S:457.1nm; and the metastable magnesium atoms can be chemically generated. A supersonic stream of Mg[³P] atoms were formed by reacting Mg[¹S] atoms with a N₂O/CO mixture. Our measurements suggest that a Mg-N₂O complex is formed and that it is subsequently reduced by reaction with CO to form metastable magnesium atoms and CO₂. The observed concentrations can be increased by optimizing the reaction conditions and significant increases (>10) are predicted based on the measured rate constant for the Mg-N₂O reaction.

CHEMILUMINESCENT REACTIONS OF METAL ATOMS WITH HALOGENS

Prof. John M. Parson
Department of Chemistry
The Ohio State University
Columbus, OH 43210

ABSTRACT

Electronic chemiluminescence has been used to probe product energy disposal in a variety of beam reactions of metals with halogen molecules. Reaction of group IIIA and IVA atoms with F_2 show highly inverted vibrational populations in the electronically excited monofluoride products. $Cu + F_2$, on the other hand, gives near statistical vibronic populations in CuF . Chemiluminescent reactions of Sn with F_2 and C_2 are greatly enhanced when Sn is generated in the metastable 1D state, and electronic state inversions are possible with SnF .

Bell Aerospace **TEXTRON**

Niagara Frontier Operations
Bell Aerospace Textron
Division of Textron Inc.

Post Office Box One
Buffalo, New York 14240
716/297-1000

ABSTRACT

TIN OXIDE AND SAMARIUM FLUORIDE STUDIES

by

J.W. Raymonda, R.J. Driscoll, T.E. Furner, W.L. Rushmore,
H.M. Thompson, and G.W. Tregay, Bell Aerospace Textron, Buffalo, NY

and

M.J. Lavan, U.S. Army Ballistic Missile Defense
Advanced Technology Center, Huntsville, Alabama

We will review the status of tin oxide and samarium fluoride as short wavelength chemical laser candidates. The kinetic and spectroscopic data bases will be summarized and Bell's methods for producing high fluxes of metal atom vapors will be described. Emission spectra, number densities and radiation rates obtained at Bell will be presented and prospects for the future will be outlined.

*Funded by U.S. Army Ballistic Missile Defense, Advanced Technology Center, Contracts DASG60-78-C-0030 and DASG60-83-C-0021.

Review of BaO/Ba Transfer Laser Studies

N. Djeu

University of South Florida

ABSTRACT

In the mid and late 70's, the concept of a BaO/Ba chemically pumped energy transfer laser was explored experimentally at the Naval Research Laboratory. The intended laser line in Ba was the ($6p\ ^1D-5d\ ^1D$) transition at $1.5\ \mu$. The upper laser level was to be pumped by energy transfer from excited states of BaO or BaF formed from chemical reactions between Ba and N_2O or NF_3 . The studies took the forms of both direct gain/loss measurements at $1.5\ \mu$ in a Ba + N_2O flame and relaxation measurements of the lower laser level in a number of potential diluent gases. A review of the results from these investigations will be given.

POTENTIAL OF METAL/OXIDIZER SYSTEMS

Prof. J. L. Gole
School of Physics
Georgia Institute of Technology
Atlanta, GA 30332

ABSTRACT

The potential use of metal oxidizer systems to produce chemically driven visible laser candidates is reviewed. A few basic parameters and criteria on which one would like to focus in producing either a directly driven chemical system or a pre-chemically driven-energy transfer laser system are outlined and exemplified. Direct chemically driven systems should be considered which are characterized by selective excited state formation commensurate with high quantum yields and excited state radiative lifetimes on the order of 10^{-6} to 10^{-4} seconds. Previous quantum yield determinations will be critically reviewed both as they (1) apply to direct excited state formation and (2) influence plausible energy transfers to produce lasing media. The criteria for selectivity and high quantum yield require activation energies for excited state formation less than or equal to those for ground state formation and the temperature dependence for any metathesis producing the excited states of interest must be considered. We also consider the possible use of collision induced ultrafast energy transfer ($\sigma_{\text{Trans.}} \gg \text{gas kinetic}$) among the excited states of high temperature molecules to produce population inversions. Finally we emphasize the possible use of various analogs of the $\text{O}_2(^1\Delta) - \text{I}_2$ energy transfer laser system.

WHY SO FEW CHEMICAL LASERS?

David L. Huestis
Chemical Physics Laboratory
SRI International
Menlo Park, CA 94025

ABSTRACT

A variety of flames are known for their efficiency in producing visible emission. These formed the impetus if not the basis for the search for a visible chemical laser, which has been thus far unsuccessful. Here we present a qualitative theoretical analysis of the electronic potential energy surfaces accessible to the reaction complex and a semiquantitative determination of the probability of diabatic passage through the avoided crossings. We conclude that except for spin-conserving reactions of light atoms, ground state reactants should preferentially yield ground state products. Thus the search for a visible chemical laser should emphasize [1] identification of classes of reactions with high yields of excited electronic states (rather than high yields of fluorescence) and [2] investigation of the mechanisms of chemical reactions that could be used to convert the stored electronic energy into a suitable upper laser level.

ABSTRACT: Short Wavelength Chemical Laser Workshop

Charleston, SC

14, 15 November 1984

HF LASERS - WHAT HAVE WE LEARNED?

Walter R. Warren, Jr.

Pacific Applied Research

6 Crestwind Drive

Rancho Palos Verdes, CA 90274

Hydrogen fluoride (and deuterium fluoride) vibration/rotation chemical lasers, because of their potentials in several high energy laser applications, have benefited from a considerable amount of R and D activity during the past 15 years, and a large body of knowledge now exists on the phenomena governing their operation and growth potentials. Milestones in the development of HF lasers are summarized by discussing, first, contributions from the disciplines of physical chemistry, fluid dynamics, optics, and systems requirements, and second, several interactions between these disciplines that have resulted in important advances. Attempts are made to identify lessons learned that should aid in the development of short wavelength chemical lasers.

ELECTRONIC STATE ENERGY TRANSFER AND LASER POSSIBILITIES

D. W. Setser
Kansas State University
Manhattan, KS 66506

Some excitation-transfer and dissociative excitation-transfer reactions of $N_2(A)$ will be reviewed. There is considerable evidence that quenching with formation of the Franck-Condon favored vibrational states of $N_2(X, v'')$ is the most favorable pathway. Recent studies by the flowing-afterglow technique of excitation-transfer reactions from $N_2(A)$ to SO and to S_2 fit this general model. The total quenching rate constants are large and the product branching fractions for formation of $SO(A^3\Pi)$ and $S_2(B^3\Sigma_u^-)$ appear to be favorable, especially for the SO reaction. In the second part of the presentation, work in our laboratory to develop a suitable flowing-afterglow source for kinetic and spectroscopic studies of $NF(b)$ will be discussed. A summary of some total deactivation rate constants for $NF(b)$ will be presented and compared to the quenching rate constants for $O_2(b)$ and $NH(b)$.

THE EXCITATION OF THE IF ($B^3\Pi_0^+$) STATE BY ELECTRONIC ENERGY
TRANSFER FROM $N_2(A^3\Sigma_u^+)$ *

Lawrence G. Piper, William J. Marinelli, B. David Green
and Wilson T. Rawlins

Physical Sciences Inc., Andover, MA 01810

We have studied the electronic energy transfer reaction between $N_2(A^3\Sigma_u^+, v'=0,1)$ and $IF(X^1\Sigma^+)$. We find strong excitation of the $IF(B^3\Pi_0^+)$ state with a rate coefficient of $1.2 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. At low pressures, all vibrational levels of $IF(B)$ up through $v'=8$ are about equally populated. At pressures of a few torr, the populations of the $IF(B)$ shift to the lower vibrational levels, indicating vibrational relaxation of the B-state by argon atoms. The rate coefficient for the quenching of $N_2(A)$ by $IF(X)$ is $2.0 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. Thus about 60% of the quenching interactions between $N_2(A)$ and $IF(X)$ result in $IF(B)$ excitation.

* Supported by the Air Force Weapons Laboratories under Contract
#F29601-83-C-0051

NF($b^1\Sigma^+$) -- IF TRANSFER¹

A. T. PRITT, JR. and D. J. BENARD
Rockwell International Science Center
Thousand Oaks, CA 91360 USA

Production of IF($B^3\Pi(O^+)$), observed in the $F + HN_3 + CF_3I$ (or I_2) flame, has been attributed to a near-resonant electronic energy transfer from NF($b^1\Sigma^+$) to IF($X^1\Sigma^+$).² This mechanism has been confirmed by generating a transient concentration of NF($b^1\Sigma^+$) in the presence of ground state IF.

The E-E energy transfer rate constant and the observed vibrational distribution of the $B^3\Pi(O^+)$ state depend on the flame stoichiometry. The results of these experiments are explained in terms of an electronic energy transfer mechanism requiring that the individual state-to-state rates are proportional to the probabilities associated with "vertical" transitions.

¹This work supported under Air Force Contract F29601-83-C-0048

²A. T. Pritt, Jr, D. Patel, and D. J. Benard, Chem Phys Lett. **97**, 471 (1983).

NO(A \rightarrow X) Laser
M. D. Burrows
Los Alamos National Laboratory

Abstract

Lasing action on the (0,1) and (0,2) γ -bands on NO at 237 and 248 nm, respectively, has been obtained by longitudinal pumping of 0.5 Torr of NO on the (0,0) γ -band transition at 227 nm. Typical output pulse energies were 13 μ J at an intrinsic energy conversion efficiency of approximately 15%. Lasing is observed from single rotational lines with linewidths limited only by the NO room temperature Doppler width of 0.1 cm^{-1} . The potential for lasing on other NO γ -bands will also be discussed. The (2,0) band at 205 nm can be pumped by a Raman shifted (first stokes line from D₂, narrow linewidth ArF laser. The resulting NO laser would then operate on the ($v' = 2, 1 \leq v'' \leq 8$) bands in the 214 to 220 nm wavelength region. If sufficiently fast relaxation within the vibrational manifold of the A² Σ state can be achieved, lasing on the ($v' = 1, 1 \leq v'' \leq 7$) bands in the 225 to 290 nm region should also be possible.

Vibration-to-Electronic Energy
Transfer in V-V Pumped Molecules

J. W. Rich
Calspan Advanced Technology Center
P. O. Box 400
Buffalo, New York 14225

Extreme nonequilibrium overpopulation of high energy vibrational states can be produced in a variety of molecules by vibration-to-vibration (V-V) energy exchange pumping. Recent work is reviewed which shows substantial population of vibrational states above 4 eV in translationally cold CO, N₂, and NO by this mechanism. The process is operative at high, collision-dominated densities, can occur with a variety of means of vibrational mode excitation (electrical, optical, or chemical reaction), and can occur in either pulsed or steady-state experiments.

Current studies show that energy can be transferred from such V-V pumped vibrational levels to selected electronic state acceptors. The CO a ³Π and A ¹Π states are observed to be populated by transfer from V-V pumped CO X ¹Σ⁺; NO A ²Σ⁺ and B ²Π levels are similarly populated when NO X ²Π is V-V pumped. Intermolecular energy transfer has also been observed, notably transfer from CO X ¹Σ⁺ to the NO B ²Π, NO A ²Σ⁺, and HgBr B ²Σ states. The application of such vibration-to-electronic energy transfer processes to short wavelength laser action is discussed.

"Electronic Energy Transfer in Metal Atoms"

Stephen R. Leone*

Joint Institute for Laboratory Astrophysics
National Bureau of Standards and University of Colorado
Boulder, Colorado 80309

A number of ideas for producing short wavelength gas lasers and excimer lasers involve storage of energy in excited states of atoms, followed by collisional or radiative transfer of this energy to the lasing states. Electronic excitation transfer is thus of fundamental importance to the viability and efficiency of such schemes. Detailed results will be presented on state-to-state energy transfer pathways, cross sections, and transfer efficiencies for both Na systems (work of A. Gallagher, JILA) and Ca systems (Leone). These investigations elucidate some of the major factors involved in near resonant energy transfer, collisional up-pumping, associative ionization, and spin-changing collisions. The results provide insight into necessary propensity rules for these energy transfer processes. Recent experiments demonstrate that orbital "alignment" can have a significant influence on near resonant energy transfer processes. These experiments can be a valuable tool in elucidating mechanisms for the major energy transfer pathways in these systems.

*Staff member, Quantum Physics Division, National Bureau of Standards.

Abstract of a technical presentation at the
SHORT WAVELENGTH CHEMICAL LASER WORKSHOP
held in Charleston, South Carolina, November 14 & 15, 1984

Thomas F. George
Department of Chemistry
University of Rochester
Rochester, New York 14627
(716)275-3586

Hai-Woong Lee
Department of Physics
Oakland University
Rochester, Michigan 48063
(313)377-3419

PRODUCTION OF EXCITED ATOMIC STATES BY ION-SURFACE CHARGE EXCHANGE

The intensity of radiation emitted from ion-surface charge exchange processes can be enhanced if the surface exposed to impinging ions is electronically excited. Semiclassical calculations are carried out for the probabilities of electron transfer for Li^{3+} , alpha particles (He^{2+}) and He^+ colliding with a Si(111) surface, where a laser is used to excite electrons in silicon from the valence band to surface states. It is shown that with a moderate-power laser ($\sim \text{W}/\text{cm}^2$), high inversion densities of Li^{2+} , He^+ and He can be obtained as necessary for high gain.

Inelastic and Reactive Collisions of Electronically Excited Species

Millard H. Alexander
Department of Chemistry
University of Maryland
College Park MD 20742

Paul J. Dagdigian
Department of Chemistry
The Johns Hopkins University
Baltimore MD 21218

An overview will be given of some new experimental techniques for the study of inelastic and reactive collisions of atoms and molecules in low-lying excited states, with emphasis on spin-orbit as well as rotational state specificity. The current level of understanding of the dynamics of these processes will also be described.

ATTENDANCE LIST

Dr Hal Ahlstrom
Lawrence Livermore Natl Lab
PO BOX 808
Livermore CA 94550

Prof. M. H. Alexander
Dept. of Chemistry
University of Maryland
College Park MD 20742

Dr A.A. Alfano
Rockwell Science Center
PO BOX 1085
Thousand Oaks CA 91360

Dr. P. Avizonis
Kirtland Air Force Base
AFWL/AR
Albuquerque NM 87117

Dr R.W. Bass
Litton Guidance & Control Syst
5500 Canoga Ave MS 67/35
Woodland Hills CA 91365

Dr J Batteh
Science Applications Inc
1503 Johnson Ferry, #100
Marietta GA 30062

Mr. S.L. Baughcum
Los Alamos Natl.
PO BOX 1663, CHM-4 MS J567
Los Alamos N.M. 87545

Dr. D. Benard
Rockwell Science Center
P.O. Box 1085
Thousand Oaks CA 91360

Dr. J. Betts
TRW
One Space Park, R1 1184
Redondo Beach CA 90278

Dr G Black
SRI Intl, PN-019
333 Ravenswood Ave
Menlo Park CA 94025

Dr Jay Blauer
Rocketdyne, FA44
6633 Canoga Avenue
Canoga Park CA 91304

Dr. R. Burnham
Naval Research Lab
Code 6540
Washington DC 20375

Prof. N. Djeu
Univ. South Florida
Department of Physics
Tampa FL 33620

Dr. M. Burrows
Los Alamos National Lab
PO BOX 1663, MS E 543
Los Alamos NM 87545

Mr T Dreiling
The Boeing Company
PO BOX 3707
Seattle WA 98124

Dr Geo Bush
KMS Fusion
PO BOX 1567
Ann Arbor MI 48106

Mr R Driscoll
Bell Aerospace Textron
PO BOX 1
Buffalo NY 14240

Dr Geo Caledonia
Physical Sciences Inc
Research Park, PO BOX 3100
Andover MA 01810

Dr M Duignan
Potomac Photonics Inc
2233 Arlington Terr
Alexandria VA 22303

Dr R Carmen
Rocketdyne, FA 44
6633 Canoga Ave
Canoga Park CA 91304

Dr. D. J. Eckstrom
Stanford Research Institute
Menlo Park CA 94025

Dr. E. Chicklis
Sanders Associates Inc.
95 Canal St.
Nashua NH 03061

Prof. G. Eden
Dept. of Electrical Engr.
University of Illinois
Urbana IL 61801

Mr Chas Clendening
TRW R1 1162
One Space Park
Redondo Beach CA 90278

Mr D English
TRW, R1 1184
One Space Park
Redondo Beach CA 90278

Prof. T. Cool
Dept. of Applied & Engr. Phys.
Cornell University
Ithaca NY 14853

Dr. J. J. Ewing
Mathematical Sciences NW, Inc.
2755 Northrup Way
Bellevue WA 98004

Prof. R. D. Coombe
Department of Chemistry
University of Denver
Denver CO 80208

Dr. W. Felder
Aerochem Research Laboratories
Princeton NJ 08540

Prof. P. Davidovits
Department of Chemistry
Boston College
Chestnut Hill MA 02167

Dr. B. Feldman
Naval Research Laboratory
Code 6540
Washington DC 20375

Dr. S. Davis
AFWL/ARDD
Kirtland AFB
Albuquerque NM 87117

Prof. A. Fontijn
Dept. of Chemical Engineering
R. P. I.
Troy NY 12180

Prof. Jack Gelfand
Princeton University
D-420 Engineering School
Princeton NJ 08544

Dr J Herbelin
Aerospace Corporation
PO BOX 91055
Los Angeles CA 90009

Dr. T. F. George
Department of Chemistry
U. Rochester,
Rochester NY 14627

Dr. J. Hinchin
UTRC
Silver Lane
East Hartford CT 06108

Dr. A. Glass
President, KMS Fusion Inc.
Box 1567
Ann Arbor MI 48106

Dr. J. F. Hon
Rocketdyne, FA44
6633 Canoga Avenue
Canoga Park CA 91304

Dr D A Gobeli
Rockwell Science Center
PO BOX 1085
Thousand Oaks CA 91360

Dr A.B. Horowitz
TRW
One Space Park
Redondo Beach CA 90278

Prof. J. Gole
Department of Chemistry
Georgia Tech.
Atlanta GA 30332

Dr J Hough
The Aerospace Corporation
PO BOX 91005
Los Angeles CA 90009

Dr. R. Greiner
Los Alamos National Lab
PO BOX 1663, MS G 740
Los Alamos NM 87545

Dr S Howie
Science Applications Inc
1710 Goodridge Dr.
McLean VA 22102

Dr. R. Guenther
US Army Research Office
PO BOX 12211
Res Triangle Pk NC 27709

Dr. D. L. Huestis
Stanford Research Institute
Menlo Park CA 94025

Mr F Hanson
NOSC
271 Catalina Blvd
San Diego CA 92152

Dr. A. Hyder
Auburn University
202 Samford Hall
Auburn AL 36849

Dr Geo Harpole
TRW, R1 1184
One Space Park
Redondo Beach CA 90278

Dr. T. A. Jacobs
Assoc Deputy Assist Secy Navy
Pentagon, Rm. 4D-745
Washington DC 20350

Mr A Hartford
Los Alamos Natl Lab, CHM-DO
PO BOX 1663
Los Alamos NM 87545

Dr W Jeffers
Helios Inc
1822 Sunset Place
Longmont CO 80501

Dr. R. Heidner
The Aerospace Corporation
P.O. Box 92957, M5741
Los Angeles CA 90009

Dr. R.C. Jensen
Hercules Aerospace Division
Bacchus Works
Magna UT 84044

Dr. C.R. Jones
Los Alamos National Lab
PO BOX 1663, MS J564
Los Alamos NM 87545

Dr. M. Lavan
BMDATC/T
PO BOX 1500
Huntsville AL 35807

Dr P Jones
Dept of Chem. Ohio State Univ
140 W 18th Ave
Columbus OH 43210

Prof J Lawless
Dept of Chemistry
Carnegie Mellon Univ
Pittsburg PA 15213

Dr. R. Junker
ONR Code 412
800 N Quincy Street
Arlington VA 22203

Dr. C. Leader
McDonnell Douglas Research Lab
P.O. Box 576
St. Louis MO 63166

Prof. F. Kaufman
University of Pittsburgh
Dept of Chemistry
Pittsburgh PA 15261

Prof H.W. Lee
Dept of Physics
Oakland University
Rochester MI 48036

Dr. J. D. Kelley
McDonnell Douglas Research Lab
PO BOX 576
St. Louis MO 63166

Dr. S. R. Leone
JILA Campus Box 440
University of Colorado
Boulder CO 80309

Dr. R.L. Kerber
Divn Engineering Research
Michigan State University
East Lansing MI 48824

Dr. D.H. Leslie
W.J.Schafer Associates Inc
1901 N Fort Myer Dr
Arlington VA 22209

Dr P Kleiber
Iowa Laser Facility
University of Iowa
Iowa City IA 52242

Dr. M. J. Linevsky
Applied Physics Lab.
Johns Hopkins University
Laurel MD 20810

LtCol D. Kline
SDIO/OSD
Washington DC 20301

Dr R Lontz
US Army Research Office
PO BOX 12211
Res Triangle Pk NC 27709

Dr J.B. Koffend
The Aerospace Corporation
PO BOX 91055
Los Angeles CA 90009

Capt Doug Loverro
AFWL/ARA
Kirtland AFB
Albuquerque NM 87117

Dr R.C. Kulander
Lawrence Livermore Natl Lab
PO BOX 5508
Livermore CA 94550

Mr Bob Malins
BDM Corporation
1801 Randolph Rd., S.E.
Albuquerque NM 87106

Mr Herb Lager
Rocketdyne, FA 44
6633 Canoga Avenue
Canoga Park CA 91304

Dr. L. Marquet
SDIO/OSD
Washington DC 20301

Dr Paul Marrone
Calspan Adv Technology Center
PO BOX 400
Buffalo NY 14225

Dr Jacob Maya
GTE Sylvania
100 Endicott
Danvers MA 01923

Dr. R. Meinzer
UTRC
Silver Lane
East Hartford CT 06108

Dr. H. Michels
UTRC
Silver Lane
East Hartford CT 06108

Capt. L. Myers
AFOSR/NC
Bolling AFB
Washington DC 20332

Dr. James Newton
FTD-TQTD
Wright Patterson AFB
OH 45433

Dr. W.L. Nikolai
Hercules Aerospace Division
Bacchus Works
Magna UT 84044

Dr. R. Oldenborg
Los Alamos National Laboratory
P.O. Box 1663, MS J567
Los Alamos NM 87545

Dr. L. Palumbo
W.J.Schafer Associates Inc
Corporate Place 128,Bld 2
Wakefield MA 01880

Dr. J. Parsons
Chem.Dept.,140 W. 18th Ave
Ohio State University
Columbus OH 43210

Dr. K. Patterson
W.J.Schafer Associates Inc.
1901 N Fort Myer Dr
Arlington VA 22209

Mr N Pchelkin
AFWL/ARAY
Kirtland AFB
Albuquerque NM 87117

Mr W.K. Pendelton
AFWL/AR-1
Kirtland AFB
Albuquerque NM 87117

Dr Otis Peterson
Los Alamos National Lab
PO BOX 1663,MS E527
Los Alamos NM 87545

Dr H Pilloff
ONR Code 412
800 N Quincy St
Arlington VA 22203

Dr. L. Piper
Physical Sciences Inc.
Research Park,PO BOX 3100
Andover MA 01810

Mr D Plummer
RDA
321 Hermosa Dr SE
Albuquerque NM 87108

Dr. H. Powell
Lawrence Livermore Natl Lab
PO BOX 808, MS L-490
Livermore CA 94550

Dr A.T. Pritt
Rockwell Science Center
PO BOX 1085
Thousand Oaks CA 91360

Prof H Rabitz
Dept of Chemistry
Princeton University
Princeton NJ 08544

Dr. J. Raymonda
Bell Aerospace Textron
P.O. Box 1
Buffalo NY 14240

Dr. L. Redmon
Chemical Dynamics
1550 W. Henderson Rd
Columbus OH 43220

Dr. K. Reed
Sanders Associates Inc
95 Canal Street
Nashua NH 03061

Prof. D. Setser
Kansas State University
Manhattan Kansas 66502

Dr T.N. Rescigno
Lawrence Livermore Natl Lab
PO BOX 808
Livermore CA 94550

Dr R.W. Shaw
US Army Research Office
PO BOX 12211
Res Triangle Pk NC 27709

Dr J. Rice
Sandia National Laboratory
Dept 1120
Albuquerque NM 87185

Dr R Slater
Avco Everett Research Lab
2385 Revere Beach Pkwy
Everett MA 02149

Dr. J. W. Rich
Calspan Corporation
4455 Genesee Street
Buffalo NY 14221

Mr W Smith
Science Applications Inc
1503 Johnson Ferry Rd 100
Marietta GA 30062

Capt Mark Rogers
AFWL/AR
Kirtland AFB
Albuquerque NM 87117

Dr. W. R. Sooy
Lawrence Livermore Lab, L-488
P.O. Box 808
Livermore CA 94550

Dr. M. Rossi
Stanford Research Institute
Menlo Park CA 94025

Dr Alan Stanton
Aerodyne Research Inc
45 Manning Rd
Billerica MA 01821

Capt Ted Salvi
AFWL/ARBF
Kirtland AFB
Albuquerque NM 87117

Dr. D. H. Stedman
Department of Chemistry
University of Denver
Denver CO 80208

Dr E Schimitschek
NOSC Code 811
271 Catalina Blvd
San Diego CA 92152

Dr. D. Sutton
The Aerospace Corporation
P.O. Box 92957, M5741
Los Angeles CA 90009

Dr Peter Schulz
School of Physics
Georgia Technical Inst
Atlanta GA 30332

Maj Orv Swensen
AFWL/ARDD
Kirtland AFB
Albuquerque NM 87117

Dr. S. K. Searles
Naval Research Lab
Code 6540
Washington DC 20375

Dr J.C. Swingle
Lawrence Livermore Natl Lab
PO BOX 808, MS L-6
Livermore CA 94550

Prof. L. Sentman
Aeronautical & Astro Engr.
University of Illinois
Urbana IL 61801

Dr Hung Tai-Wang
Naval Research Laboratory
Code 6180
Washington DC 20375

Dr. R. L. Taylor
CVD Incorporated
35 Industrial Parkway
Woburn MA 01801

Prof. J. Weiner
Chemistry Department
University of Maryland
College Park MD 20742

Dr. D.L. Thompson
Oklahoma State University
Dept of Chemistry
Stillwater OK 74078

Prof. J. Weisenfeld
Department of Chemistry
Cornell University
Ithaca NY 14853

Dr W.P. Thompson
The Aerospace Corporation
PO BOX 92957, MS M5/742
Los Angeles CA 90009

Dr. M. White
Off. of Naval Rsrch Detachment
495 Summer St, Barnes Bld
Boston MA 02210

Dr Geo Tregay
Bell Aerospace Textron
PO BOX 1
Buffalo NY 14240

Prof. C. Wittig
Univ. Southern California
Department of Chemistry
Los Angeles CA 90089

Dr Jack Trost
TRW
One Space Park
Redondo Beach CA 90278

Mr Ron Yates
Dow Chemical Co
Bldg 1776
Midland MI 48640

Dr C Ultee
United Technologies Res Ctr
Silver Lane
E. Hartford CT 06108

Dr C Von Rosenberg
Avco Everett Research Lab
2385 Revere Beach Pkwy
Everett MA 02149

Dr. W. Warren
Pacific Applied Rsrch. Corp.
6 Crestwind Drive
Rcho Palos Vdes CA 90274

Prof W.S. Warren
Dept of Chemistry
Princeton University
Princeton NJ 08544

Dr. W.S. Watt
W.J.Schafer Associates Inc
1901 N Fort Myer Dr
Arlington VA 22209

Dr. J. Waymouth
GTE Sylvania
100 Endicott Street
Danvers MA 01923